

Genmab to Showcase Latest Blood Cancer Treatment Advancements at 2025 American Society of Hematology (ASH) Annual Meeting

Media Release

COPENHAGEN, Denmark; November 3, 2025

- More than 20 abstracts, including seven oral presentations, highlight advancements from the epcoritamab development program supporting the expanded clinical profile and potential of epcoritamab for a broader range of B-cell malignancies
- New data demonstrate potential of epcoritamab in first and second-line settings in follicular lymphoma (FL) and first-line in diffuse large B-cell lymphoma (DLBCL)
- Results from Phase 3 EPCORE FL-1 trial, evaluating epcoritamab in combination with rituximab and lenalidomide (R²) in patients with relapsed or refractory (R/R) follicular lymphoma (FL) accepted for oral presentation

Genmab A/S (Nasdaq: **GMAB**) today announced that more than 20 abstracts evaluating epcoritamab-bysp, a T-cell engaging bispecific antibody administered subcutaneously, across lines of therapy and B-cell non-Hodgkin's lymphoma (NHL) subtypes, will be presented at the 67th Annual Meeting and Exposition of the American Society of Hematology (ASH), in Orlando, Florida, and online, December 6-9.

Data from the epcoritamab development program will showcase its expanding clinical profile and potential utility in earlier lines of therapy with a fixed treatment duration. Presentations include three oral sessions supporting the potential of epcoritamab in the first- and second-line setting in patients with follicular lymphoma (FL) and two oral presentations evaluating epcoritamab in the first-line setting in patients with diffuse large B-cell lymphoma (DLBCL). Additionally, two oral presentations will summarize the efficacy and safety of epcoritamab as monotherapy and in combination for patients with Richter transformation (RT).

"The breadth and depth of data evaluating epcoritamab at this year's American Society of Hematology meeting spotlight the growing body of clinical evidence supporting the potential of epcoritamab and underscore our commitment to developing epcoritamab as a potential core therapy across a range of B-cell malignancies," said Dr. Judith Klimovsky, Executive Vice President and Chief Development Officer of Genmab. "We look forward to sharing our data at ASH, including the full pivotal results from the Phase 3 EPCORE FL-1 trial evaluating epcoritamab in combination with rituximab and lenalidomide in patients with relapsed or refractory follicular lymphoma."

2025 R&D Update and ASH Data Review

On Thursday, December 11 at 11:00 a.m. ET/5:00 p.m. CEST, Genmab will host its 2025 R&D Update and ASH Data Review. The event will be virtual and webcast live. Details, including the webcast link and registration will be available on www.genmab.com. This meeting is not an official program of the ASH Annual Meeting.

All abstracts accepted for presentation have been published and may be accessed on the [ASH website](https://www.aschm.org). The following abstracts evaluating epcoritamab have been accepted for presentation at ASH:

Oral Presentations

Abstract Number	Abstract Title	Type of Presentation	Date/Time of Presentation
63	Vitolo et al., Fixed-duration epcoritamab monotherapy induces high response and MRD-negativity rates in elderly patients with newly diagnosed large B-cell lymphoma (LBCL) and comorbidities: results from EPCORE DLBCL-3	Oral	December 6, 9:30 - 11:00 AM (Presentation: 10:00 AM - 10:15 AM)

Genmab to Showcase Latest Blood Cancer Treatment Advancements at 2025 American Society of Hematology (ASH) Annual Meeting

Abstract Number	Abstract Title	Type of Presentation	Date/Time of Presentation
64	Cheah et al., Epcoritamab + R-mini-CHOP results in 2-year remissions and high MRD negativity rates in elderly patients with newly diagnosed DLBCL: results from the EPCORE NHL-2 trial	Oral	December 6, 9:30 - 11:00 AM (Presentation: 10:15 - 10:30 AM)
464*	Merryman et al., Rituximab and epcoritamab as first-line therapy for patients with high-tumor burden follicular lymphoma: Results of a multicenter phase II trial	Oral	December 7, 9:30-11:00 AM (Presentation: 9:45 - 10:00 AM)
465	Leslie et al., Epcoritamab with rituximab + lenalidomide (R ²) and epcoritamab maintenance deliver deep and durable remissions in previously untreated (1L) follicular lymphoma (FL): 3-year outcomes from EPCORE NHL-2 arms 6 and 7	Oral	December 7, 9:30 - 11:00 AM (Presentation: 10:00 - 10:15 AM)
466	Falchi et al., Primary phase 3 results from the EPCORE FL-1 trial of epcoritamab with rituximab and lenalidomide (R ²) versus R ² for relapsed or refractory follicular lymphoma	Oral	December 7, 9:30 AM - 11:00 AM (Presentation: 10:15 - 10:30 AM)
1015	Thompson et al., Epcoritamab combinations demonstrate promising efficacy in patients (pts) with Richter transformation (RT): first results from arms 2B (epcor + lenalidomide [LEN]) and 2C (epcor + R-CHOP) of the phase 1b/2 EPCORE CLL-1 trial	Oral	December 8, 4:30 - 6:00 PM (Presentation: 4:30 - 4:45 PM)
1017	Kater et al., Epcoritamab monotherapy demonstrates promising efficacy in patients with Richter transformation (RT): 2-year follow-up results from arm 2A of the phase 1b/2 EPCORE CLL-1 trial	Oral	December 8, 4:30 - 6:00 PM (Presentation: 5:00 - 5:15 PM)

*Investigator-led trial

Poster Presentations

Abstract Number	Abstract Title	Type of Presentation	Date/Time of Presentation
1820	Noorani et al., Optimal dose of epcoritamab in combination with lenalidomide and rituximab in relapsed or refractory follicular lymphoma – analysis of pharmacokinetics and exposure-response relationships of EPCORE FL-1 phase 3 study	Poster	December 6, 5:30 - 7:30 PM
1955	Falchi et al., Fixed-duration epcoritamab + R-CHOP in patients with newly diagnosed DLBCL and high IPI scores (3-5) led to sustained remissions and disease-free survival beyond 3-years: results from the EPCORE NHL-2 trial	Poster	December 6, 5:30 - 7:30 PM
1959	Torres Lopez et al., Outpatient administration of epcoritamab monotherapy for relapsed/refractory diffuse large B-cell lymphoma (R/R DLBCL): results from the EPCORE NHL-6 by race and ethnicity	Poster	December 6, 5:30-7:30 PM

Genmab to Showcase Latest Blood Cancer Treatment Advancements at 2025 American Society of Hematology (ASH) Annual Meeting

Abstract Number	Abstract Title	Type of Presentation	Date/Time of Presentation
1960	Thieblemont et al., Epcoritamab (epcore) monotherapy offers long-term disease control in large B-cell lymphoma (LBCL): NHL-1 subgroup analysis in patients with prior chimeric antigen receptor T-cell (CAR T) therapy from the 3-year follow-up	Poster	December 6, 5:30 - 7:30 PM
2721	Park et al., Barriers to receiving CAR T-cell treatment among patients with non-Hodgkin lymphoma who were deemed eligible for CAR T-cell therapy	Poster	December 6, 5:30 - 7:30 PM
3565	Robinson et al., Phenotype and functional state of endogenous T-cells support T-cell engager therapy in the post-CAR T setting	Poster	December 7, 6:00 - 8:00 PM
3566	Takacs et al., Exposure to epcoritamab is associated with improved T-cell functionality and dynamic changes in CD8+ T-cells in diffuse large B-cell lymphoma: insights from EPCORE NHL-6	Poster	December 7, 6:00 - 8:00 PM
3736	Brody et al., Epcoritamab + GemOx achieves durable >2-year remissions in relapsed/refractory (R/R) 2L+ diffuse large B-cell lymphoma (DLBCL): long-term data reinforce clinical potential of the regimen across a diverse patient population	Poster	December 7, 6:00 - 8:30 PM
4481	Xavier et al., Underreporting of prognostic factors in real-world studies for bispecifics in relapsed or refractory diffuse large B-cell lymphoma	Poster	December 7, 6:00 - 8:00 PM
5511	Cheah et al., Durable responses in patients with large B-cell lymphoma and 3+ prior lines of therapy who either paused or discontinued epcoritamab monotherapy while in complete response	Poster	December 8, 6:00 - 8:00 PM
5513	Karimi et al., Sustained remissions beyond 4 years with epcoritamab monotherapy: long-term follow-up results from the pivotal EPCORE NHL-1 trial in patients with relapsed or refractory large B-cell lymphoma	Poster	December 8, 6:00 - 8:00 PM
5357	Vitolo et al., Fixed-duration epcoritamab in combination with bendamustine + rituximab (BR) for first-line (1L) treatment of follicular lymphoma (FL): 3-year results from EPCORE NHL-2 arm 3 demonstrate deep and durable responses with manageable safety	Poster	December 8, 6:00 - 8:00 PM
5370	Linton et al., HRQoL in relapsed/refractory follicular lymphoma patients treated with epcoritamab in combination with rituximab plus lenalidomide (E+R ²): primary results of patient-reported outcomes from the EPCORE FL-1 trial	Poster	December 8, 6:00 - 8:00 PM
5393	Strati et al., EPCORE FL-2 phase 3 trial of epcoritamab with rituximab and lanalidomide (R ²) vs chemoimmunotherapy (CIT) in previously untreated follicular lymphoma (FL): trial in progress	Poster	December 8, 6:00 - 8:00 PM

Genmab to Showcase Latest Blood Cancer Treatment Advancements at 2025 American Society of Hematology (ASH) Annual Meeting

e-Publications

Abstract Number	Abstract Title	Type of Presentation	Date/Time of Presentation
7251	Johnson et al., Epcoritamab monotherapy provides superior efficacy vs non-anthracycline-containing regimens in newly diagnosed elderly DLBCL patients deemed unsuitable for anthracycline-containing regimens: a match-adjusted comparative efficacy analysis	Publication	NA
7942	Graff et al., Operational efficiencies and cost savings of using one bispecific antibody FDA-approved for both R/R 3L+ DLBCL and FL	Publication	NA
8075	Ali et al., Effectiveness of epcoritamab in a heterogeneous population with relapsed/refractory diffuse large B-cell lymphoma including post-chimeric antigen receptor T-cell therapy patients: insights from the real-world epcoritamab patient characteristics and outcomes research (Real-EPCOR) study	Publication	NA

The safety and efficacy of epcoritamab have not been established for these investigational uses.

About Epcoritamab

Epcoritamab is an IgG1-bispecific antibody created using Genmab's proprietary DuoBody® technology and administered subcutaneously. Genmab's DuoBody-CD3 technology is designed to direct cytotoxic T cells selectively to elicit an immune response toward target cell types. Epcoritamab is designed to simultaneously bind to CD3 on T cells and CD20 on B cells and induces T-cell-mediated killing of CD20+ cells.¹

Epcoritamab (approved under the brand name EPKINLY® in the U.S. and Japan, and TEPKINLY® in the EU) has received regulatory approval in certain lymphoma indications in several territories. Epcoritamab is being co-developed by Genmab and AbbVie as part of the companies' oncology collaboration. The companies will share commercial responsibilities in the U.S. and Japan, with AbbVie responsible for further global commercialization. Both companies will pursue additional international regulatory approvals for the investigational R/R FL indication and additional approvals for the R/R DLBCL indication.

Genmab and AbbVie continue to evaluate the use of epcoritamab as a monotherapy, and in combination, across lines of therapy in a range of hematologic malignancies. This includes five ongoing Phase 3, open-label, randomized trials including a trial evaluating epcoritamab as a monotherapy in patients with R/R DLBCL compared to investigators choice chemotherapy ([NCT04628494](#)), a trial evaluating epcoritamab in combination with R-CHOP in adult patients with newly diagnosed DLBCL ([NCT05578976](#)), a trial evaluating epcoritamab in combination with rituximab and lenalidomide (R²) in patients with R/R FL ([NCT05409066](#)), a trial evaluating epcoritamab in combination with rituximab and lenalidomide (R²) compared to chemoimmunotherapy in patients with previously untreated FL ([NCT06191744](#)), and a trial evaluating epcoritamab in combination with lenalidomide compared to chemotherapy infusion in patients with R/R DLBCL ([NCT06508658](#)). The safety and efficacy of epcoritamab has not been established for these investigational uses. Please visit www.clinicaltrials.gov for more information.

Genmab to Showcase Latest Blood Cancer Treatment Advancements at 2025 American Society of Hematology (ASH) Annual Meeting

About Genmab

Genmab is an international biotechnology company with a core purpose of guiding its unstoppable team to strive toward improving the lives of patients with innovative and differentiated antibody therapeutics. For 25 years, its passionate, innovative and collaborative team has invented next-generation antibody technology platforms and leveraged translational, quantitative and data sciences, resulting in a proprietary pipeline including bispecific T-cell engagers, antibody-drug conjugates, next-generation immune checkpoint modulators and effector function-enhanced antibodies. By 2030, Genmab's vision is to transform the lives of people with cancer and other serious diseases with knock-your-socks-off (KYSO) antibody medicines.®

Established in 1999, Genmab is headquartered in Copenhagen, Denmark, with international presence across North America, Europe and Asia Pacific. For more information, please visit [Genmab.com](https://www.genmab.com) and follow us on [LinkedIn](#) and [X](#).

Contact:

David Freundel, Senior Director, Global Communications & Corporate Affairs

T: +1 609 613 0504; E: dafr@genmab.com

Andrew Carlsen, Vice President, Head of Investor Relations

T: +45 3377 9558; E: acn@genmab.com

This Media Release contains forward-looking statements. The words "believe," "expect," "anticipate," "intend" and "plan" and similar expressions identify forward looking statements. Actual results or performance may differ materially from any future results or performance expressed or implied by such statements. The important factors that could cause our actual results or performance to differ materially include, among others, risks associated with preclinical and clinical development of products, uncertainties related to the outcome and conduct of clinical trials including unforeseen safety issues, uncertainties related to product manufacturing, the lack of market acceptance of our products, our inability to manage growth, the competitive environment in relation to our business area and markets, our inability to attract and retain suitably qualified personnel, the unenforceability or lack of protection of our patents and proprietary rights, our relationships with affiliated entities, changes and developments in technology which may render our products or technologies obsolete, and other factors. For a further discussion of these risks, please refer to the risk management sections in Genmab's most recent financial reports, which are available on www.genmab.com and the risk factors included in Genmab's most recent Annual Report on Form 20-F and other filings with the U.S. Securities and Exchange Commission (SEC), which are available at www.sec.gov. Genmab does not undertake any obligation to update or revise forward looking statements in this Media Release nor to confirm such statements to reflect subsequent events or circumstances after the date made or in relation to actual results, unless required by law.

Genmab A/S and/or its subsidiaries own the following trademarks: Genmab®; the Y-shaped Genmab logo®; Genmab in combination with the Y-shaped Genmab logo®; HuMax®; DuoBody®; HexaBody®; DuoHexaBody®, HexElect® and KYSO®. EPCORE®, EPKINLY®, TEPKINLY® and their designs are trademarks of AbbVie Biotechnology Ltd.

¹ Engelberts PJ, Hiemstra IH, de Jong B, et al. DuoBody-CD3xCD20 induces potent T-cell-mediated killing of malignant B cells in preclinical models and provides opportunities for subcutaneous dosing. *EBioMedicine*. 2020;52:102625. DOI: 10.1016/j.ebiom.2019.102625.