

European Commission approves label expansion of Roche's Hemlibra to include people with moderate haemophilia A in the EU

- **Hemlibra, already approved for severe haemophilia A in the EU, will now also provide an effective and convenient prophylactic treatment option for people with moderate haemophilia A**
- **Moderate haemophilia A can have a significant impact on the lives of people affected, with only 15% living a bleed-free life¹**
- **The approval is based on the HAVEN 6 results, where Hemlibra demonstrated effective bleed control and a favourable safety profile in people with moderate haemophilia A without inhibitors²**

Basel, 01 February 2023 - Roche (SIX: RO, ROG; OTCQX: RHHBY) today announced that the European Commission approved the expansion of the Hemlibra® (emicizumab) European Union (EU) marketing authorisation. The label will now include the routine prophylaxis of bleeding episodes in people with haemophilia A (congenital factor VIII deficiency) without factor VIII inhibitors, who have moderate disease (FVIII $\geq 1\%$ and $\leq 5\%$) with a severe bleeding phenotype.³ Haemophilia A affects around 900,000 people worldwide,^{4,5} approximately 14% of whom have a moderate form of the disease.⁶

“We welcome the European Commission's decision to approve Hemlibra also for people with moderate haemophilia A in the EU, since even moderate disease can produce bleeds that cause irreversible joint damage and impact quality of life,” said Levi Garraway, M.D., Ph.D., Roche's Chief Medical Officer and Head of Global Product Development. “As its benefit expands to broader populations, we remain committed to determining how Hemlibra might help even more people with haemophilia A to live a bleed-free life.”

While the treatment and care of people with severe haemophilia A are well-established, there is less information and guidance on prophylaxis for moderate haemophilia A.⁷ Additionally, the severity of haemophilia A, traditionally measured by an individual's factor VIII levels, is not always reflective of bleeding behaviour. Many people with moderate haemophilia A may not be receiving prophylactic treatments and could have a worsened clinical burden. Approximately 85% of people with moderate haemophilia A have bleeds within a given year and one in three have long-term joint problems, many of which require surgery and impact quality of life.^{1,7,8}

This approval is based on the results of the phase III HAVEN 6 trial, in which Hemlibra demonstrated effective bleed control and a favourable safety profile in people with non-severe haemophilia A without factor VIII inhibitors, where prophylaxis was clinically indicated.² The decision was also based on real world data. This label expansion will provide

an effective and convenient prophylactic treatment option with a favourable safety profile for people in the EU with moderate haemophilia A with a severe bleeding phenotype.

Hemlibra is approved as a prophylactic treatment option for people with haemophilia A with factor VIII inhibitors in more than 110 countries and for people without factor VIII inhibitors in more than 100 countries worldwide. It has been studied in one of the largest clinical trial programmes in people with haemophilia A with and without factor VIII inhibitors, including eight phase III studies.

About HAVEN 6

HAVEN 6 is a phase III clinical study designed to evaluate the safety, efficacy, pharmacokinetics and pharmacodynamics of Hemlibra in people with non-severe haemophilia A without factor VIII inhibitors. Data from the primary analysis of HAVEN 6 was presented at the 30th International Society on Thrombosis and Haemostasis (ISTH) Annual Congress, on 11 July 2022.² The analysis included data from 72 participants, including three women, 51 (70.8%) of whom had moderate haemophilia A without factor VIII inhibitors. The data indicate that Hemlibra has effective bleed control and a favourable safety profile in people with non-severe haemophilia A without factor VIII inhibitors, with no new safety signals identified. Hemlibra achieved clinically meaningful bleed control, with 66.7% of participants experiencing no bleeds that required treatment, 81.9% experiencing no spontaneous bleeds that required treatment, and 88.9% experiencing no joint bleeds that required treatment. Model-based annualised bleed rates (ABR) remained low throughout the evaluation period at 0.9 (95% CI: 0.55-1.52).²

About Hemlibra® (emicizumab)

Hemlibra is a bispecific factor IXa- and factor X-directed antibody. It is designed to bring together factor IXa and factor X, proteins involved in the natural coagulation cascade, and restore the blood clotting process for people with haemophilia A. Hemlibra is a prophylactic (preventative) treatment that can be administered by an injection of a ready-to-use solution under the skin (subcutaneously) once-weekly, every two weeks, or every four weeks (after an initial once-weekly dose for the first four weeks). Hemlibra was created by Chugai Pharmaceutical Co., Ltd. and is being co-developed globally by Chugai, Roche and Genentech. It is marketed in the United States by Genentech as Hemlibra (emicizumab-kxwh), with kxwh as the suffix designated in accordance with Nonproprietary Naming of Biological Products Guidance for Industry issued by the U.S. Food and Drug Administration.⁹

About haemophilia A

Haemophilia A is an inherited, serious disorder in which a person's blood does not clot properly, leading to uncontrolled and often spontaneous bleeding. Haemophilia A affects around 900,000 people worldwide,^{4,5} approximately 14% of whom have a moderate form of the disorder.⁶ However, the severity of haemophilia A is not always reflective of bleeding behaviour, as some people with non-severe haemophilia may experience symptoms similar to

those with severe haemophilia and warrant prophylaxis.⁸ All severities of haemophilia A can significantly reduce the quality of life for people affected, as well as their family and caregivers. People with haemophilia A either lack or do not have enough of a clotting protein called factor VIII. In a healthy person, when a bleed occurs, factor VIII brings together the clotting factors IXa and X, which is a critical step in the formation of a blood clot to help stop bleeding. Depending on the severity of their symptoms, people with haemophilia A can bleed frequently, especially into their joints or muscles.⁶ These bleeds can present a significant health concern as they often cause pain and can lead to chronic swelling, deformity, reduced mobility and long-term joint damage.¹⁰ A serious complication of treatment is the development of inhibitors to factor VIII replacement therapies. Inhibitors are antibodies developed by the body's immune system that bind to and block the efficacy of replacement factor VIII, making it difficult, if not impossible, to obtain a level of factor VIII sufficient to control bleeding.¹¹

About Roche in haematology

Roche has been developing medicines for people with malignant and non-malignant blood diseases for more than 20 years; our experience and knowledge in this therapeutic area runs deep. Today, we are investing more than ever in our effort to bring innovative treatment options to patients across a wide range of haematological diseases. Our approved medicines include MabThera[®]/Rituxan[®] (rituximab), Gazyva[®]/Gazyvaro[®] (obinutuzumab), Polivy[®] (polatuzumab vedotin), Venclexta[®]/Venclyxto[®] (venetoclax) in collaboration with AbbVie, Hemlibra[®] (emicizumab) and Lunsumio[®] (mosunetuzumab). Our pipeline of investigational haematology medicines includes T-cell engaging bispecific antibodies glofitamab, targeting both CD20 and CD3, and cevostamab, targeting both FcRH5 and CD3; Tecentriq[®] (atezolizumab), a monoclonal antibody designed to bind with PD-L1 and crovalimab, an anti-C5 antibody engineered to optimise complement inhibition. Our scientific expertise, combined with the breadth of our portfolio and pipeline, also provides a unique opportunity to develop combination regimens that aim to improve the lives of patients even further.

About Roche

Founded in 1896 in Basel, Switzerland, as one of the first industrial manufacturers of branded medicines, Roche has grown into the world's largest biotechnology company and the global leader in in-vitro diagnostics. The company pursues scientific excellence to discover and develop medicines and diagnostics for improving and saving the lives of people around the world. We are a pioneer in personalised healthcare and want to further transform how healthcare is delivered to have an even greater impact. To provide the best care for each person we partner with many stakeholders and combine our strengths in Diagnostics and Pharma with data insights from the clinical practice.

In recognising our endeavour to pursue a long-term perspective in all we do, Roche has been named one of the most sustainable companies in the pharmaceuticals industry by the Dow

Jones Sustainability Indices for the thirteenth consecutive year. This distinction also reflects our efforts to improve access to healthcare together with local partners in every country we work.

Genentech, in the United States, is a wholly owned member of the Roche Group. Roche is the majority shareholder in Chugai Pharmaceutical, Japan.

For more information, please visit www.roche.com.

All trademarks used or mentioned in this release are protected by law.

References

- [1] Nissen F, et al. An Insight into clinical outcomes in mild, moderate, and severe hemophilia A (HA): A preliminary analysis of the CHES II study. Presented at: International Society on Thrombosis and Haemostasis (ISTH) Virtual Congress; 2020, July 12-14. Abstract OC 09.3.
- [2] Hermans C, et al. Efficacy of Emicizumab Prophylaxis for the Treatment of People with Moderate or Mild Hemophilia A without Factor VIII Inhibitors: Results from the Primary Analysis of the HAVEN 6 Study. Presented at: International Society on Thrombosis and Haemostasis (ISTH) Congress; 2022, July 11. Abstract OC 30.5.
- [3] Hemlibra, INN-emicizumab sMPC [Internet; cited 2023 January]. Available from: <https://ec.europa.eu/health/documents/community-register/html/h1271.htm>
- [4] Srivastava A, et al. WFH Guidelines for the Management of Hemophilia, 3rd edition. Haemophilia. 2020; 26 (Suppl 6): 1- 158.
- [5] Iorio A, et al. Establishing the Prevalence and Prevalence at Birth of Hemophilia in Males. Ann Intern Med 2019 Oct 15;171(8):540-546.
- [6] World Federation of Hemophilia. Report on the annual global survey 2020. [Internet; cited 2023 January] Available from: <http://www1.wfh.org/publications/files/pdf-2045.pdf>.
- [7] Walsh C, et al. Identified unmet needs and proposed solutions in mild-to-moderate haemophilia: A summary of opinions from a roundtable of haemophilia experts. Haemophilia. 2021;27(S1):25-32.
- [8] Collins PW, et al. Clinical phenotype of severe and moderate haemophilia: Who should receive prophylaxis and what is the target trough level? Haemophilia, 2021;27: 192-198.
- [9] Hemlibra, emicizumab-kxwh. U.S. Food and Drug Administration Prescribing Information [Internet; cited 2023 January]. Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/761083s000lbl.pdf
- [10] Franchini M, et al. Haemophilia A in the third millennium. Blood Rev. 2013;27(4):179-84.
- [11] Srivastava A, et al. WFH Guidelines for the Management of Hemophilia, 3rd edition. Haemophilia. 2020; 26 (Suppl 6): 1- 158.

Roche Group Media Relations

Phone: +41 61 688 8888 / e-mail: media.relations@roche.com

Hans Trees, PhD

Phone: +41 79 407 72 58

Nathalie Altermatt

Phone: +41 79 771 05 25

Karsten Kleine

Phone: +41 79 461 86 83

Nina Mählitz

Phone: +41 79 327 54 74

Dr. Barbara von Schnurbein

Phone: +41 79 699 97 44

Sileia Urech

Phone: +41 79 935 81 48

Roche Investor Relations

Dr. Bruno Eschli

Phone: +41 61 68-75284

e-mail: bruno.eschli@roche.com

Dr. Sabine Borngräber

Phone: +41 61 68-88027

e-mail: sabine.borngraeber@roche.com

Dr. Birgit Masjost

Phone: +41 61 68-84814

e-mail: birgit.masjost@roche.com

Dr. Gerard Tobin

Phone: +41 61 68-72942

e-mail: gerard.tobin@roche.com

Investor Relations North America

Loren Kalm

Phone: +1 650 225 3217

e-mail: kalm.loren@gene.com