



## Media Release

### January 31, 2023

## Idorsia submits European marketing authorisation application for aprocitentan for the treatment of patients with resistant hypertension

- The application includes data from the Phase 3 registration study of patients with resistant hypertension, where aprocitentan demonstrated a sustained blood pressure reduction over 48 weeks and was well-tolerated

### **Allschwil, Switzerland – January 31, 2023**

Idorsia Ltd (SIX: IDIA) today announced that it has submitted a marketing authorisation application (MAA) to the European Medicines Agency (EMA) for aprocitentan, Idorsia's investigational, novel dual endothelin receptor antagonist, for the treatment of patients with resistant hypertension.

The MAA includes data from a comprehensive clinical and non-clinical development program. In the Phase 3 registration study, PRECISION, aprocitentan showed statistically significant and clinically meaningful reduction in blood pressure (BP) which was maintained for up to 48 weeks when added to combination background antihypertensive therapy in patients with resistant hypertension. In PRECISION, aprocitentan was generally well tolerated with no major safety concerns. The most frequent adverse event with aprocitentan was mild-to-moderate edema/fluid retention.

Full results from the PRECISION study were published in November 2022 in *The Lancet* "[A randomized controlled trial of the dual endothelin antagonist aprocitentan for resistant hypertension](#)". More details and commentary can be found in the dedicated [press release](#) and an [investor webcast](#) featuring Prof. Markus Schlaich, an investigator in PRECISION, which are available on Idorsia's corporate website: [www.idorsia.com](http://www.idorsia.com)

Patients with uncontrolled blood pressure are at risk of major cardiovascular events.<sup>1</sup> These risks are even higher for patients whose blood pressure is uncontrolled despite treatment with three or more antihypertensives<sup>2</sup>, known as resistant hypertension<sup>3,4</sup>. It has been more than 30 years since a new anti-hypertensive therapy working by a new mechanism has been brought to patients. By targeting a currently unopposed pathophysiologic pathway, aprocitentan represents a potential novel, effective, and well-tolerated treatment for resistant hypertension.

Idorsia is developing aprocitentan together with Janssen Biotech, Inc., one of the Janssen Pharmaceutical Companies of Johnson & Johnson. Janssen Biotech has sole commercialization rights worldwide.

---

### **Notes to the editor**

#### **The endothelin system in systemic hypertension**

Endothelin-1 (ET-1) is a potent vasoconstrictor that also induces neurohormonal activation, vascular hypertrophy and remodeling, cardiac hypertrophy and fibrosis, and endothelial dysfunction. In hypertension, both ET<sub>A</sub> and ET<sub>B</sub> receptors mediate harmful effects of ET-1.<sup>5</sup> As a vasoconstrictor, co-mitogenic agent, linking pulse pressure and vascular remodeling, and mediator of aldosterone and catecholamine release, endothelin is a key player in hypertension and end-organ damage.<sup>6,7</sup>

### About resistant hypertension

Hypertension (high blood pressure) is one of the most common cardiovascular risk factors, and its prevalence continues to rise. According to a recent study, there are more than 1.3 billion people living with hypertension worldwide<sup>8</sup> – a startling number, which has almost doubled in the past 40 years. Left uncontrolled, people have a greater risk of life-threatening conditions such as heart attack, stroke, and chronic kidney disease.<sup>9</sup>

Patients with hypertension can often successfully control their blood pressure by combining a healthier lifestyle with effective medication. However, approximately 10% of patients have resistant hypertension where the blood pressure remains high despite receiving at least three antihypertensive medications of different pharmacological classes, including a diuretic, at optimal doses,<sup>3,10</sup>

The endothelin pathway has been implicated in the pathogenesis of hypertension, especially in volume- and salt-dependent forms, which are a common feature in patients with resistant hypertension, but it is currently not targeted therapeutically, thereby leaving this relevant pathophysiologic pathway unopposed with currently available medications.<sup>3,11,12</sup> This pathway is activated in patients prone to developing resistant hypertension, such as Black or African American patients, patients with obesity or obstructive sleep apnea,<sup>13-15</sup> and in comorbid conditions frequently associated with resistant hypertension such as diabetes and chronic kidney disease.<sup>16-19</sup>

### About aprocitentan

Aprocitentan is an investigational, novel, oral, dual endothelin receptor antagonist (ERA), which potently inhibits the binding of ET-1 to ET<sub>A</sub> and ET<sub>B</sub> receptors. Aprocitentan has a low potential for drug-drug interaction and a mechanism of action that is ideally suited for the pathophysiology of resistant hypertension.

### About PRECISION<sup>20,21</sup> (NCT03541174)

PRECISION was a multicenter, blinded, randomized, parallel-group, Phase 3 study, which was performed in hospitals or research centers in Europe, North America, Asia, and Australia. Patients were eligible for randomization if their sitting systolic blood pressure was 140 mm Hg or higher despite taking standardized background therapy consisting of three antihypertensive drugs, including a diuretic. The study consisted of three sequential parts: Part 1 was the 4-week double-blind, randomized, and placebo-controlled part, in which 730 patients were randomized to aprocitentan 12.5 mg (n=243), aprocitentan 25 mg (n=243), or placebo (n=244) in a 1:1:1 ratio; Part 2 was a 32-week single (patient)-blind part, in which all patients received aprocitentan 25 mg (n=704); and Part 3 was a 12-week double-blind, randomized, and placebo-controlled withdrawal part, in which patients were re-randomized to aprocitentan 25 mg (n=307) or placebo (n=307) in a 1:1 ratio. The primary and key secondary endpoints were changes in unattended office systolic blood pressure from baseline to week 4 and from withdrawal baseline to week 40, respectively. Secondary endpoints included 24-h ambulatory blood pressure changes.

At baseline, 69.2% of patients were obese or severely obese, 54.1% had diabetes, 22.2% had stage 3-4 chronic kidney disease and 19.6% had congestive heart failure. At screening, 63% of all patients who were randomly assigned were prescribed four or more antihypertensive drugs.

### Key PRECISION findings<sup>21</sup>

The least square mean change in office SBP at 4 weeks was –15.3 mmHg for aprocitentan 12.5 mg, –15.2 mmHg for 25 mg, and –11.5 mmHg for placebo, for a difference versus placebo of **–3.8 mmHg** (p=0.0042) and **–3.7 mmHg** (p=0.0046), respectively. Office diastolic blood pressure (DBP) also decreased with both aprocitentan doses compared to placebo (–3.9 mmHg for the 12.5 mg dose and –4.5 mmHg for the 25 mg dose). Office SBP and DBP were maintained during Part 2 in patients previously receiving aprocitentan and decreased within the first 2 weeks of Part 2 before stabilizing in those previously receiving placebo. In Part 3, office SBP after 4 weeks of withdrawal (the key secondary endpoint) increased significantly with placebo compared to aprocitentan (**5.8 mmHg**; p<0.0001). Office DBP also increased with placebo compared to aprocitentan (5.2 mmHg; p<0.001). The difference between the two groups remained up to week 48.

The results from ambulatory BP monitoring confirmed those derived from office measurements. At the end of Part 1, aprocitentan, after placebo correction, decreased both the 24-hour ambulatory SBP (**–4.2 mmHg for the 12.5 mg dose** and **–5.9 mmHg for the 25 mg dose**) and DBP (–4.3 mmHg for the 12.5 mg dose and –5.8 mmHg for the 25 mg dose). The placebo-corrected SBP lowering effect was –5.1 mmHg and –7.4 mmHg during the nighttime and –3.8 mmHg and –5.3 mmHg during the daytime, for the 12.5 mg and 25 mg doses, respectively. In Part 3, after 4 weeks of withdrawal (week 40), both the 24-hour ambulatory SBP and DBP increased with placebo compared with aprocitentan (6.5 mm Hg and 6.8 mm Hg respectively).

Treatment-emergent adverse events (TEAEs) during the 4-week double-blind study period (Part 1) were reported in 27.6% and 36.7% of the patients treated with 12.5 and 25 mg aprocitentan, respectively, versus 19.4% in the placebo group. The most frequent adverse event was fluid retention which was reported more frequently with aprocitentan than with placebo in a dose-dependent fashion (9.1%, 18.4%, and 2.1% for patients receiving aprocitentan 12.5 mg, 25 mg and placebo, during Part 1, respectively; 18.2% for patients receiving aprocitentan 25 mg during Part 2; and 2.6% and 1.3% for patients on aprocitentan 25 mg and placebo, during Part 3, respectively). Fluid retention was generally mild-to-moderate. Discontinuation due to edema/fluid retention was reported for seven patients.

### About the collaboration agreement with Janssen Biotech, Inc.

In 2017, Idorsia entered into a collaboration agreement with Janssen Biotech, Inc., one of the Janssen Pharmaceutical Companies of Johnson & Johnson, to jointly develop aprocitentan and any of its derivative compounds or products. Idorsia received a one-time milestone payment of USD 230 million. Both parties have joint development rights over aprocitentan. Idorsia has conducted the Phase 3 development and is overseeing the regulatory review for the treatment of patients with difficult-to-control hypertension. The costs are shared equally between both partners. Janssen Biotech, Inc. has sole commercialization rights worldwide, whereas Idorsia is entitled to receive tiered royalties on annual net sales in each calendar year (20% up to USD 500 million, 30% from USD 500 million up to USD 2.0 billion, and 35% above USD 2.0 billion) for the licensed products in the collaboration indications. Janssen Biotech, Inc. will oversee the Phase 3 development and submission for any additional indications.

### Key Literature

1. Turnbull F. Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-designed overviews of randomized trials. *Lancet* 2003; 362: 1527–35
2. Kumbhani DJ, et al. Resistant hypertension: a frequent and ominous finding among hypertensive patients with atherothrombosis. *European Heart Journal* (2013) 34, 1204–1214.
3. Williams B, et al. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J* 2018; 39: 3021–104.
4. Carey RM, et al. Resistant Hypertension: Detection, Evaluation, and Management A Scientific Statement From the American Heart Association. *Hypertension*. 2018;72:00-00.
5. Dolan E, et al. Superiority of ambulatory over clinic blood pressure measurement in predicting mortality: the Dublin outcome study. *Hypertension* 2005; 46:156–61.
6. Kedzierski RM, et al. Endothelin system: the double-edged sword in health and disease. *Annu Rev Pharmacol Toxicol*. 2001; 41:851-76.
7. Iglarz M, et al. At the heart of tissue: endothelin system and end-organ damage. *Clin Sci* 2010; 119:453-63
8. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet* 2021; 398:957-80.
9. Daugherty SL, et al. Incidence and prognosis of resistant hypertension in hypertensive patients. *Circulation*. 2012 Apr 3;125(13):1635-42.
10. Noubiap JJ, et al. Global prevalence of resistant hypertension: a meta-analysis of data from 3.2 million patients. *Heart* 2019; 105: 98–105.
11. Dhaun N, et al. Role of endothelin-1 in clinical hypertension: 20 years on. *Hypertension* 2008; 52:452-9.
12. Clozel M. Aprocitentan and the endothelin system in resistant hypertension. *Can J Physiol Pharmacol* 2022; 100:573-83.
13. Grubbs AL, et al. Saphenous vein endothelin system expression and activity in African American patients. *Arterioscler Thromb Vasc Biol* 2002; 22: 1122–7.
14. Parrinello G, et al. Central obesity and hypertension: the role of plasma endothelin. *Am J Hypertens* 1996; 9: 1186–91.
15. Phillips BG, et al. Effects of obstructive sleep apnea on endothelin-1 and blood pressure. *J Hypertens* 1999; 17: 61–6.
16. Takahashi K, et al. Elevated plasma endothelin in patients with diabetes mellitus. *Diabetologia* 1990; 33: 306–10.
17. Solini A, et al. Resistant hypertension in patients with type 2 diabetes: clinical correlates and association with complications. *J Hypertens* 2014; 32: 2401–10; discussion 10.
18. Dhaun N, Webb DJ, Kluth DC. Endothelin-1 and the kidney--beyond BP. *Br J Pharmacol* 2012; 167: 720–31.
19. Rossignol P, et al. The double challenge of resistant hypertension and chronic kidney disease. *Lancet* 2015; 386: 1588–98.
20. Danaïetash P et al. Identifying and treating resistant hypertension in PRECISION: A randomized long-term clinical trial with aprocitentan. *J Clin Hypertension* 2022 Jul;24(7):804-813.
21. Schlaich MP, et al. A randomized controlled trial of the dual endothelin antagonist aprocitentan for resistant hypertension. *The Lancet*, 2022; Available online 7 November 2022



### About Idorsia

Idorsia Ltd is reaching out for more – We have more ideas, we see more opportunities and we want to help more patients. In order to achieve this, we will develop Idorsia into a leading biopharmaceutical company, with a strong scientific core.

Headquartered near Basel, Switzerland – a European biotech-hub – Idorsia is specialized in the discovery, development, and commercialization of small molecules to transform the horizon of therapeutic options. Idorsia has a broad portfolio of innovative drugs in the pipeline, an experienced team of professionals covering all disciplines from bench to bedside, state-of-the-art facilities, and a strong balance sheet – the ideal constellation to translate R&D efforts into business success.

Idorsia was listed on the SIX Swiss Exchange (ticker symbol: IDIA) in June 2017 and has over 1'200 highly qualified specialists dedicated to realizing our ambitious targets.

### For further information, please contact

Andrew C. Weiss

Senior Vice President, Head of Investor Relations & Corporate Communications

Idorsia Pharmaceuticals Ltd, Hegenheimermattweg 91, CH-4123 Allschwil

+41 58 844 10 10

[investor.relations@idorsia.com](mailto:investor.relations@idorsia.com) | [media.relations@idorsia.com](mailto:media.relations@idorsia.com) | [www.idorsia.com](http://www.idorsia.com)

The above information contains certain "forward-looking statements", relating to the company's business, which can be identified by the use of forward-looking terminology such as "estimates", "believes", "expects", "may", "are expected to", "will", "will continue", "should", "would be", "seeks", "pending" or "anticipates" or similar expressions, or by discussions of strategy, plans or intentions. Such statements include descriptions of the company's investment and research and development programs and anticipated expenditures in connection therewith, descriptions of new products expected to be introduced by the company and anticipated customer demand for such products and products in the company's existing portfolio. Such statements reflect the current views of the company with respect to future events and are subject to certain risks, uncertainties and assumptions. Many factors could cause the actual results, performance or achievements of the company to be materially different from any future results, performances or achievements that may be expressed or implied by such forward-looking statements. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those described herein as anticipated, believed, estimated or expected.